P.02

KPR-T3229A-RE

Serial No. 10/655,988

Markings show differences between previously issued claims and newly allowed reissue claims.

A polymer composition, produced by the steps of: reacting one or more 1. compounds with the terminal functional groups on a polymer, and said polymer consists essentially of:

polymerized 1,3-butadiene having a peak molecular weight between 500 and 20,000, 1,2addition between 30% and 70%, and hydrogenation of at least 90% of the unsaturation; a ratio of viscosity (in poise at room temperature) to peak molecular weight raised to the 3.4 power of at most 2.0 times 10.9; and

[one]about two or more terminal functional groups per molecule.

- The polymer composition of claim [1]6, wherein the terminal functional groups 2. of the polymer are selected from [a]the group consisting of hydroxyl, carboxyl, phenol, epoxy, and amine groups.
- Canceled. 3.
- The polymer composition of claim [3]6, wherein the polymerized 1,3-butadiene 4. has a peak molecular weight between 1,000 and 10,000.
- The polymer composition of claim 4, wherein the polymerized 1,3-butadiene is at 5. least 95% hydrogenated.
- The polymer composition of claim [5]1, wherein the ratio of viscosity to peak 6. molecular weight raised to the 3.4 power of the polymer is less than 1.0 [x]times 10-9.
- The polymer composition of claim 6, wherein the terminal functional groups of 7. the polymer consist of about two hydroxyl groups per molecule.

- 8. The polymer composition of claim 1, wherein the peak molecular weight of the polymer is between 1000 and 10000.
- 9. The polymer composition of claim [8]6, wherein the 1,2-addition of the polymerized 1.3-butadiene is between 40% and 60%.
- 10. The polymer composition of claim 1, wherein the polymerized 1,3-butadiene has about two hydroxyl groups per molecule.
- 11. The [polymeric]polymer composition of claim 10, wherein the polymerized 1,3-butadiene is reacted with compounds that form a coating.
- 12. The [polymeric]polymer composition of claim 10, wherein the polymerized 1,3-butadiene is reacted with compounds that form a block selected from the group consisting of polyesters, polyamides, and polycarbonates.
- 13. The polymer composition of claim 8 wherein the polymer has a peak molecular weight of about 10,000.
- 14. The polymer composition of claim 8 wherein the polymer has a peak molecular weight of about 5,000.
- 15. The polymer composition of claim 8 wherein the polymer has a peak molecular weight of about 3,000.
- 16. The polymer composition of claim 8 wherein the polymer has a peak molecular weight of about 2,000.
- 17. The polymer composition of claim 4 wherein the polymer has a peak molecular weight of about 10,000.

P. 04

- The polymer composition of claim 4 wherein the polymer has a peak molecular weight of about 5,000.
- The polymer composition of claim 4 wherein the polymer has a peak molecular weight of about 3,000.
- The polymer composition of claim 4 wherein the polymer has a peak molecular weight of about 2,000.

## Claims 21-23 Cancelled.

- The polymer composition of claim 1 wherein the polymerized 1,3-butadiene has 1.7 terminal functional groups per molecule.
- The polymer composition of claim I wherein the polymerized 1,3-butadiene has 1.9 terminal functional groups per molecule.